IN THE CLAIMS

1. (currently amended) A compound, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound having the general structure shown in Formula I:

$$\mathbb{R}^4$$
 \mathbb{R}^3
 \mathbb{R}^1

Formula I

wherein:

Y is selected from the group consisting of the following moieties: alkyl, alkyl-aryl, heteroalkyl, heteroaryl, arylheteroaryl, alkyl-heteroaryl, cycloalkyl, alkyloxy, alkylaryloxy, aryloxy, heteroaryloxy, heterocycloalkyloxy, cycloalkyloxy, alkylamino, arylamino, alkyl-arylamino, arylamino, heteroarylamino, cycloalkylamino and heterocycloalkylamino, with the proviso that Y maybe optionally substituted with X^{11} or X^{12} ;

 X^{11} is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl, heterocyclyl, heterocyclylalkyl, aryl, alkylaryl, arylalkyl, heteroaryl, alkylheteroaryl, or heteroarylalkyl, with the proviso that X^{11} may be additionally optionally substituted with X^{12} ;

 X^{12} is hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, carboxy, carbalkoxy, carboxamido, alkoxycarbonylamino, alkoxycarbonyloxy, alkylureido, arylureido, halogen, cyano, or nitro, with the proviso that said alkyl, alkoxy, and aryl may be additionally optionally substituted with moieties independently selected from X^{12} ;

 R^1 is COR^5 or $B(OR)_2$, wherein R^5 is H, OH, OR^8 , NR^9R^{10} , CF_3 , C_2F_5 , C_2F_7 , CF_2R^6 , R^6 , or COR^7 wherein R^7 is H, OH, OR^8 , CHR^9R^{10} , or NR^9R^{10} , wherein R^6 , R^8 , R^9 and R^{10} are independently selected from the group consisting of H, alkyl, aryl, heteroalkyl, heteroaryl, cycloalkyl, cycloalkyl, arylalkyl, heteroarylalkyl, $[CH(R^{1'})]_pCOOR^{11}$, $[CH(R^{1'})]_pCONR^{12}R^{13}$, $[CH(R^{1'})]_pSO_2R^{11}$, $[CH(R^{1'})]_pCOR^{11}$, $[CH(R^{1'})]_pCH(OH)R^{11}$, $CH(R^{1'})CONHCH(R^{2'})COOR^{11}$, $CH(R^{1'})CONHCH(R^{2'})CONR^{12}R^{13}$, $CH(R^{1'})CONHCH(R^{2'})R'$, $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})COOR^{11}$, CH $(R^{1'})$ CONHCH $(R^{2'})$ CONHCH $(R^{3'})$ CONR¹²R¹³, $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONHCH(R^{4'})COOR^{11}$, CH $(R^{1'})$ CONHCH $(R^{2'})$ CONHCH $(R^{3'})$ CONHCH $(R^{4'})$ CONR¹²R¹³, $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONHCH(R^{4'})CONHCH(R^{5'})COO R^{11}$ and $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONHCH(R^{4'})CONHCH(R^{5'})CONR^{12}R^{13}$ wherein $R^{1'}$, $R^{2'}$, $R^{3'}$, $R^{4'}$, $R^{5'}$, R^{11} , R^{12} , R^{13} , and R' are independently selected from the group consisting of H, alkyl, aryl, heteroalkyl, heteroaryl, cycloalkyl, alkylaryl, alkyl-heteroaryl, aryl-alkyl and heteroaralkyl;

Z is selected from O, N, CH or CR;

- W maybe present or absent, and if W is present, W is selected from C=O, C=S, C(=N-CN), or SO_2 ;
- Q maybe present or absent, and when Q is present, Q is CH, N, P, $(CH_2)_p$, $(CHR)_p$, $(CRR')_p$, O, NR, S, or SO_2 ; and when Q is

absent, M may be present or absent; when Q and M are absent, A is directly linked to L;

A is O, CH_2 , $(CHR)_p$, $(CHR-CHR')_p$, $(CRR')_p$, NR, S, or SO_2 or a bond;

E is CH, N, CR, or a double bond towards A, L or G;

- G may be present or absent, and when G is present, G is $(CH_2)_p$, $(CHR)_p$, or $(CRR')_p$; and when G is absent, J is present and E is directly connected to the carbon atom in Formula I as G is linked to;
- J maybe present or absent, and when J is present, J is $(CH_2)_p$, $(CHR)_p$, or $(CRR')_p$, SO_2 , NH, NR or O; and when J is absent, G is present and E is directly linked to N shown in Formula I as linked to J;
- L may be present or absent, and when L is present, L is CH, CR, O, S or NR; and when L is absent, then M may be present or absent; and if M is present with L being absent, then M is directly and independently linked to E, and J is directly and independently linked to E;
- M may be present or absent, and when M is present, M is O, NR, S, SO₂, (CH₂) $_p$, (CHR) $_p$ (CHR-CHR') $_p$, or (CRR') $_p$; p is a number from 0 to 6; and
- R, R', R², R³ and R⁴ are independently selected from the group consisting of H; C₁-C₁₀ alkyl; C₂-C₁₀ alkenyl; C₃-C₈ cycloalkyl; C₃-C₈ heterocycloalkyl, alkoxy, aryloxy, alkylthio, arylthio, amino, amido, ester, carboxylic acid, carbamate, urea, ketone, aldehyde, cyano, nitro, halogen; (cycloalkyl)alkyl and (heterocycloalkyl)alkyl, wherein said cycloalkyl is made of three to eight carbon atoms, and zero to six oxygen, nitrogen, sulfur, or phosphorus atoms, and said alkyl is of one to six carbon atoms; aryl; heteroaryl; alkyl-aryl; and alkyl-heteroaryl;

wherein said alkyl, heteroalkyl, alkenyl, heteroalkenyl, aryl, heteroaryl, cycloalkyl and heterocycloalkyl moieties may be optionally and chemically-suitably substituted, with said term "substituted" referring to optional and chemically-suitable substitution with one or more moieties selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, aralkyl, cycloalkyl, heterocyclic, halogen, hydroxy, thio, alkoxy, aryloxy, alkylthio, arylthio, amino, amido, ester, carboxylic acid, carbamate, urea, ketone, aldehyde, cyano, nitro, sulfonamido, sulfoxide, sulfone, sulfonyl urea, hydrazide, and hydroxamate;

further wherein said unit N-C-G-E-L-J-N represents a five-membered or six-membered cyclic ring structure with the proviso that when said unit N-C-G-E-L-J-N represents a five-membered cyclic ring structure, or when the bicyclic ring structure in Formula I comprising N, C, G, E, L, J, N, A, Q, and M represents a five-membered cyclic ring structure, then said five-membered cyclic ring structure lacks a carbonyl group as part of the cyclic ring.


- 2. (currently amended) The compound of claim 1, wherein R^1 is COR^5 , and R^5 is H, OH, $COOR^8$, or $CONR^9R^{10}$.
- 3. (original) The compound of claim 2, wherein R^1 is $COCONR^9R^{10}$, and R^9 is H, R^{10} is H, R^{14} , $[CH(R^{1'})]_pCOOR^{11}$, $[CH(R^{1'})]_pCONR^{12}R^{13}$, $[CH(R^{1'})]_pSO_2R^{11}$, $[CH(R^{1'})]_pSO_2R^{11}$, $[CH(R^{1'})]_pCOR^{11}$, $[CH(R^{1'})]_pCOR^{11}$, $[CH(R^{1'})]_pCOR^{11}$, or $[CH(R^{1'})]_pCOR^{11}$, or $[CH(R^{1'})]_pCOR^{11}$, wherein $[CH(R^{1'})]_pCOR^{11}$, alkyl, aryl, heteroalkyl, heteroaryl, cycloalkyl, alkyl-aryl, alkyl-heteroaryl, aryl-alkyl, alkenyl, alkynyl or heteroaralkyl.
- 4. (original) The compound of claim 3, wherein R^{10} is H, R^{14} , $CH(R^{1'})COOR^{11}$, $CH(R^{1'})COOR^{11}$, $CH(R^{1'})CONR^{12}R^{13}$, $CH(R^{1'})CH(R^{1'})CONR^{12}R^{13}$, $CH(R^{1'})CH(R^{1'})CH(R^{1'})SO_2R^{11}$, $CH(R^{1'})CH(R^{1'})SO_2N$

 $R^{12}R^{13}$, $CH(R^{1'})CH(R^{1'})COR^{11}$, $CH(R^{1'})CONHCH(R^{2'})COOR^{11}$, $CH(R^{1'})CONHCH(R^{2'})$ (R'), wherein $R^{1'}$ is H or alkyl, and $R^{2'}$ is phenyl, substituted phenyl, hetero atom-substituted phenyl, thiophenyl, cycloalkyl, piperidyl or pyridyl.

- 5. (original) The compound of claim 4, wherein $R^{1'}$ is H.
- 6. (original) The compound of claim 5, wherein R^{11} is H, methyl, ethyl, allyl, tert-butyl, benzyl, α -methylbenzyl, α , α -dimethylbenzyl, 1-methylcyclopropyl or 1-methylcyclopentyl;

R' is hydroxymethyl or CH₂CONR¹²R¹³;

 $R^{2'}$ is independently selected from the group consisting of:



.....

 ${
m U}^1$ and ${
m U}^2$ maybe same or different and are selected from H, F, CH2COOH, CH2COOMe, CH2CONH2, CH2CONHMe, CH2CONMe2, azido, amino, hydroxyl, substituted amino,

substituted hydroxyl;

 \mbox{U}^3 and \mbox{U}^4 maybe same or different and are selected from O and S;

 ${\tt U}^5$ is selected from the moieties consisting of alkyl sulfonyl, aryl sulfonyl, heteroalkyl sulfonyl, heteroaryl sulfonyl, alkyl carbonyl, aryl carbonyl, heteroalkyl

Docket No.: SCHERING 3.0-122

Application No.: 09/908,955


arylaminocarbonyl,

heteroarylaminocarbonyl or a combination thereof; and $NR^{12}R^{13}$ is selected from the group consisting of:

wherein ${\rm U}^6$ is H, OH, or ${\rm CH_2OH}$, and

 R^{14} is selected from the group consisting of: H, Me, Et, n-propyl, methoxy, cyclopropyl, n-butyl, 1-but-3-ynyl, benzyl, α -methylbenzyl, phenethyl, allyl, 1-but-3-enyl, OMe, cyclopropylmethyl.

7. (original) The compound of claim 2, wherein \mathbb{R}^2 is selected from the group consisting of the following moieties:



8. (original) The compound of claim 7, wherein \mathbb{R}^3 is selected from the group consisting of:

$$CH_{3} \xrightarrow{} CH_{3} \qquad CH_{3} \xrightarrow{} CH_{3} \qquad CH_{3} \xrightarrow{} CH_{3} \qquad CH_{3} \xrightarrow{} CH_{3}$$

$$CH_{3} \xrightarrow{} CH_{3} \xrightarrow{} CH_{3} \xrightarrow{} COOEt$$

$$CH_{3} \xrightarrow{} COOT$$

Docket No.: SCHERING 3.0-122

$$H_3C \stackrel{\frown}{\downarrow} 0-3$$
 $CH_3 \stackrel{\frown}{\downarrow} SBn$ $HO \stackrel{\frown}{\downarrow} CH_3$ $CH_3 \stackrel{\frown$

wherein $R^{31} = OH$ or O-alkyl; Y^{19} is selected from the following moieties:

$$\frac{1}{2}$$
 COOH $\frac{1}{2}$ CooH $\frac{1}$

and Y^{20} is selected from the following moieties:


9. (original) The compound of claim 8, wherein \mathbb{R}^3 is selected from the group consisting of the following moieties:

$$CH_{3} \xrightarrow{} CH_{3} \xrightarrow{$$

- 10. (original) The compound of claim 9, wherein Z is N and ${\ensuremath{R^4}}$ is H.
- 11. (original) The compound of claim 10, wherein W is C=O.

12. (original) The compound of claim 11, wherein Y is selected from the following moieties:

Docket No.: SCHERING 3.0-122




Docket No.: SCHERING 3.0-122


wherein:


 Y^{11} is selected from H, COOH, COOEt, OMe, Ph, OPh, NHMe, NHAc, NHPh, CH(Me)₂, 1-triazolyl, 1-imidazolyl, and NHCH₂COOH; Y^{12} is selected from H, COOH, COOMe, OMe, F, Cl, or Br; Y^{13} is selected from the following moieties:

 Y^{14} is selected from MeSO₂, Ac, Boc, iBoc, Cbz, or Alloc; Y^{15} and Y^{16} are independently selected from alkyl, aryl, heteroalkyl, and heteroaryl;

- Y^{17} is CF_3 , NO_2 , $CONH_2$, OH, $COOCH_3$, OCH_3 , OC_6H_5 , C_6H_5 , COC_6H_5 , NH_2 , or COOH; and
- Y^{18} is COOCH₃, NO₂, N(CH₃)₂, F, OCH₃, CH₂COOH, COOH, SO₂NH₂, or NHCOCH₃.
- 13. (original) The compound of claim 12, wherein Y is selected from the group consisting of:







wherein:

 $Y^{17} = CF_3$, NO_2 , $CONH_2$, OH, NH_2 , or COOH; $Y^{18} = F$, COOH,

14. (original) The compound of claim 13, wherein Y is selected from the group consisting of:

15. (original) The compound of claim 14, wherein L and M are absent, and J is directly linked to E.

- 16. (Original) The compound of claim 14, wherein L, J and M are absent and E is directly linked to N.
- 17. (original) The compound of claim 14, wherein ${\tt G}$ and ${\tt M}$ are absent.
- 18. (original) The compound of claim 14, wherein the moiety:

19. (original) The compound of claim 18, wherein structure \underline{a} is selected from the following structures:

20. (original) The compound of claim 18, wherein structure \underline{a} is:

wherein R^{20} is selected from the following structures:

21. (original) The compound of claim 18, wherein structure \underline{a} is:

wherein R^{21} and R^{22} may be the same or different and are independently selected from the following structures:


22. (original) The compound of claim 18, wherein structure \underline{a} is selected from the following structures:

Application No.: 09/908,955 Docket No.: SCHERING 3.0-122

23. (original) The compound of claim 14, wherein:

wherein Q may be present or absent, and if Q is absent, M is directly linked to A.

24. (original) The compound of claim 23, wherein structure \underline{b} is selected from the following structures:





25. (original) The compound of claim 14, wherein:

wherein G and J are independently selected from the group consisting of $(CH_2)_p$, $(CHR)_p$, $(CHR-CHR')_p$, and $(CRR')_p$; A and M are independently selected from the group consisting of O, S, SO_2 , NR, $(CH_2)_p$, $(CHR)_p$, $(CHR-CHR')_p$, and $(CRR')_p$; and Q is CH_2 , CHR, CRR', NH, NR, O, S, SO_2 , NR, $(CH_2)_p$, $(CHR)_p$, and $(CRR')_p$.

26. (original) The compound of claim 25, wherein structure \underline{c} is selected from the following structures:

Application No.: 09/908,955 Docket No.: SCHERING 3.0-122






27. (original) The compound of claim 14, wherein:

is selected from the following structures:


Docket No.: SCHERING 3.0-122

Application No.: 09/908,955



28. (original) The compound of claim 27, wherein:

is selected from the following structures:



- 29. (original) A pharmaceutical composition comprising as an active ingredient a compound of claim 1.
- 30. (currently amended) The pharmaceutical composition of claim 29 $\underline{\text{suitable}}$ for use in treating disorders associated with hepatitis C virus (HCV).
- 31. (original) The pharmaceutical composition of claim 29 additionally comprising a pharmaceutically acceptable carrier.
- 32. (original) The pharmaceutical composition of claim 31, additionally containing an antiviral agent.
- 33. (currently amended) The pharmaceutical composition of claim 32, still additionally—further containing an interferon.
- 34. (original) The pharmaceutical composition of claim 33, wherein said antiviral agent is ribavirin and said interferon is α -interferon or pegylated interferon.
- 35. (original) A method of treating disorders associated with the HCV, said method comprising administering to a patient in need of such treatment a pharmaceutical composition which comprises therapeutically effective amounts of a compound of claim 1.
- 36. (original) The method of claim 35, wherein said administration is oral or subcutaneous.

Application No.: 09/908,955 Docket No.: SCHERING 3.0-122

37. (original) The use of a compound of claim 1 for the manufacture of a medicament to treat disorders associated with the HCV.

- 38. (original) A method of preparing a pharmaceutical composition for treating the disorders associated with the HCV, said method comprising bringing into intimate contact a compound of claim 1 and a pharmaceutically acceptable carrier.
- 39. (currently amended) A compound exhibiting hepatitis C
 virus (HCV) protease inhibitory activity, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being selected from the compounds of structures listed below:

 $(R = t\text{-butyl}, X = NH_2)$ $(R = lsobutyl, X = NH_2)$ (R = t-butyl, X = OH)

(R = Trichloroethyl, X = OH)

$$(X = O^t Bu)$$

 $(X = OH)$

(X = OH)

 $(X = O^tBu)$

(X = NH₂)

(X = NHMe)

 $(X = NMe_2)$

Docket No.: SCHERING 3.0-122

 $(X = NH_2)$

(X = NMe₂) (X = NHMe) (X = OH)

(X = OH)

(X = NH₂)

 $(X = NMe_2)$

(R = t-butyl)(R = Isobutyl)

 $(X = O^tBu)$

(X = OH)

 $(X = NH_2)$

 $(X = NMe_2)$

(X = NMeOMe)

Application No.: 09/908,955

Docket No.: SCHERING 3.0-122

Application No.: 09/908,955

Docket No.: SCHERING 3.0-122

$$\begin{array}{c|c} H_3C & CH_3 \\ \hline \\ N & NH_2 \\ \hline \\ NH & CH_3 \\ \hline \\ H_3C & O \\ \hline \\ \end{array}$$

Application No.: 09/908,955 Docket No.: SCHERING 3.0-122

Application No.: 09/908,955 Docket No.: SCHERING 3.0-122

$$H_3C \underbrace{\hspace{1cm}}_{OH_3} OH$$

H₂C_{H₃}

- 40. (currently amended) A pharmaceutical composition for treating disorders associated with the hepatitis C virus (HCV), said composition comprising therapeutically effective amount of one or more compounds in claim 39 and a pharmaceutically acceptable carrier.
- 41. (original) The pharmaceutical composition of claim 40, additionally containing an antiviral agent.
- 42. (currently amended) The pharmaceutical composition of claim 41, still additionally containing an interferon or PEGpegylated-interferon alpha conjugate.
- 43. (original) The pharmaceutical composition of claim 42, wherein said antiviral agent is ribavirin and said interferon is α -interferon.
- 44. (original) A method of treatment of a hepatitis C virus associated disorder, comprising administering an effective amount of one or more compounds of claim 39.
- 45. (original) A method of modulating the activity of hepatitis C virus (HCV) protease, comprising contacting HCV protease with one or more compounds of claim 39.
- 46. (original) A method of treating, preventing, or ameliorating one or more symptoms of hepatitis C, comprising administering an effective amount of one or more compounds of claim 39.
- 47. (original) The method of claim 45, wherein the HCV protease is the NS3/NS4a protease.
- 48. (original) The method of claim 47, wherein the compound or compounds inhibit HCV NS3/NS4a protease.

Application No.: 09/908,955

49. (original) A method of modulating the processing of hepatitis C virus (HCV) polypeptide, comprising contacting a composition containing the HCV polypeptide under conditions in which the polypeptide is processed with one or more compounds of claim 39.

50. (previously presented) The compound of claim 8, wherein R^3 is:

51. (currently amended) A compound exhibiting hepatitis C virus
<a href="https://example.com/hepatitis-c-virus-hepatitis-c-v

52. (currently amended) A compound exhibiting hepatitis C virus
(HCV) protease inhibitory activity, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being the compound of structure shown below:

Application No.: 09/908,955

53. (currently amended) A compound exhibiting hepatitis C virus
(HCV) protease inhibitory activity, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being the compound of structure shown below:

54. (currently amended) A compound exhibiting hepatitis C virus
<a href="https://exhibitory.com/hepatitis-c-virus-hepatitis-

55. (currently amended) A compound exhibiting hepatitis C virus
(HCV) protease inhibitory activity, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of

Application No.: 09/908,955 Docket No.: SCHERING 3.0-122

said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being the compound of structure shown below:

(currently amended) A compound exhibiting hepatitis C virus (HCV) protease inhibitory activity, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being the compound of structure shown below:

(currently amended) A compound exhibiting hepatitis C virus (HCV) protease inhibitory activity, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being the compound of structure shown below:

Application No.: 09/908,955 Docket No.: SCHERING 3.0-122

58. (currently amended) A compound exhibiting hepatitis C virus (HCV) protease inhibitory activity, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being the compound of structure shown below:

(currently amended) A compound exhibiting hepatitis C virus (HCV) protease inhibitory activity, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being the compound of structure shown below:

(currently amended) A compound exhibiting hepatitis C virus (HCV) protease inhibitory activity, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being the compound of structure shown below:

61. (previously presented) A pharmaceutical composition comprising as an active ingredient a compound, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being selected from the following:

- 62. (previously presented) The pharmaceutical composition of claim 61, additionally containing an antiviral agent.
- 63. (currently amended) The pharmaceutical composition of claim
- 62, still additionally further containing an interferon or PEC pegylated-interferon alpha conjugate.
- 64. (previously presented) The pharmaceutical composition of claim 63, wherein said antiviral agent is ribavirin and said interferon is α -interferon.
- 65. (previously presented) A method of treating disorders associated with the HCV, said method comprising administering to a patient in need of such treatment, a pharmaceutical composition which comprises therapeutically effective amounts of a compound, including enantiomers, stereoisomers, rotamers, tautomers, racemates and prodrug of said compound, and pharmaceutically acceptable salts or solvates of said compound, or of said prodrug, said compound being selected from the following: